

TABLE 4. DEATHS AND DEATH RATES FROM TYPHOID FEVER, BY YEAR, RALEIGH, N. C., 1913-1917

Year	Deaths			Rate per 100,000 Population		
	Total	White	Colored	Total	White	Colored
1913.....	13	7	6	66.0	56.7	81.7
1914.....	12	7	5	60.5	60.0	68.2
1915.....	9	4	5	45.0	31.6	68.3
1916.....	12	3	9	59.6	23.4	123.1
1917.....	10	3	7	49.3	23.1	95.9
Totals.....	56	24	32	56.1	37.9	87.4

NOTE.—Figures for 1913-1915, inclusive, from the United States Bureau of the Census. Figures for 1916 and 1917 from the local registry office.

TABLE 5. INFANT MORTALITY, BY YEAR, RALEIGH, N. C., 1913-1917  
(Exclusive of Stillbirths)

Year	Deaths Under 1			Rate per 1000 Live Births		
	Total	White	Colored	Total	White	Colored
1913.....	77	36	41	134.4	99.2	195.2
1914.....	105	41	64	193.4	126.9	290.9
1915.....	77	35	42	148.9	104.8	229.5
1916.....	73	24	49	132.0	70.8	229.0
1917.....	80	29	51	147.3	89.8	231.8
Totals.....	412	165	247	151.0	98.1	235.9

NOTE.—Figures for deaths 1913 to 1915, inclusive, from the United States Bureau of the Census. Figures for deaths 1916 and 1917 from the local registry office.

Based on the total number of births during 1917, registered and un-registered, the rate is as follows:

	Total	White	Colored
Deaths under 1.....	80	29	51
Rate per 1000 live births.....	145.4	89.3	225.7